



# Physics A

# PHYA1

## Unit 1 Particles, Quantum Phenomena and Electricity

### Data and Formulae Booklet

#### DATA FUNDAMENTAL CONSTANTS AND VALUES

Quantity	Symbol	Value	Units
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$
charge of electron	$e$	$-1.60 \times 10^{-19}$	C
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	$\alpha$	$2.90 \times 10^{-3}$	m K
electron rest mass (equivalent to $5.5 \times 10^{-4}$ u)	$m_e$	$9.11 \times 10^{-31}$	kg
electron charge/mass ratio	$e/m_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$
proton rest mass (equivalent to 1.00728 u)	$m_p$	$1.67(3) \times 10^{-27}$	kg
proton charge/mass ratio	$e/m_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$
neutron rest mass (equivalent to 1.00867 u)	$m_n$	$1.67(5) \times 10^{-27}$	kg
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$
atomic mass unit (1u is equivalent to 931.3 MeV)	u	$1.661 \times 10^{-27}$	kg

#### GEOMETRICAL EQUATIONS

arc length	$= r\theta$
circumference of circle	$= 2\pi r$
area of circle	$= \pi r^2$
surface area of cylinder	$= 2\pi rh$
volume of cylinder	$= \pi r^2 h$
area of sphere	$= 4\pi r^2$
volume of sphere	$= \frac{4}{3}\pi r^3$

#### ASTRONOMICAL DATA

Body	Mass/kg	Mean radius/m
Sun	$1.99 \times 10^{30}$	$6.96 \times 10^8$
Earth	$5.98 \times 10^{24}$	$6.37 \times 10^6$

## AS FORMULAE

## PARTICLE PHYSICS

## Rest energy values

class	name	symbol	rest energy /MeV
photon	photon	$\gamma$	0
lepton	neutrino	$\nu_e$	0
		$\nu_\mu^\pm$	0
	electron	$e^\pm$	0.510999
		$\mu^\pm$	105.659
		$\pi^\pm$	139.576
mesons	$\pi$ meson	$\pi^0$	134.972
		$K^\pm$	493.821
		$K^0$	497.762
baryons	proton	p	938.257
	neutron	n	939.551

## Properties of quarks

antiquarks have opposite signs

type	charge	baryon number	strangeness
<b>u</b>	$+\frac{2}{3}e$	$+\frac{1}{3}$	0
<b>d</b>	$-\frac{1}{3}e$	$+\frac{1}{3}$	0
<b>s</b>	$-\frac{1}{3}e$	$+\frac{1}{3}$	-1

## Properties of Leptons

	Lepton number
particles: $e^-, \nu_e; \mu^-, \nu_\mu$	+1
antiparticles: $e^+, \bar{\nu}_e; \mu^+, \bar{\nu}_\mu$	-1

## Photons and Energy Levels

photon energy	$E = hf = hc/\lambda$
photoelectricity	$hf = \phi + E_{K(\max)}$
energy levels	$hf = E_1 - E_2$
de Broglie Wavelength	$\lambda = \frac{h}{p} = \frac{h}{mv}$

## ELECTRICITY

$$\text{current and pd} \quad I = \frac{\Delta Q}{\Delta t} \quad V = \frac{W}{Q} \quad R = \frac{V}{I}$$

$$\text{emf} \quad \varepsilon = \frac{E}{Q} \quad \varepsilon = I(R + r)$$

$$\text{resistors in series} \quad R = R_1 + R_2 + R_3 + \dots$$

$$\text{resistors in parallel} \quad \frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$$

$$\text{resistivity} \quad \rho = \frac{RA}{L}$$

$$\text{power} \quad P = VI = I^2R = \frac{V^2}{R}$$

$$\text{alternating current} \quad I_{\text{rms}} = \frac{I_0}{\sqrt{2}} \quad V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

## MECHANICS

$$\text{moments} \quad \text{moment} = Fd$$

$$\text{velocity and acceleration} \quad v = \frac{\Delta s}{\Delta t} \quad a = \frac{\Delta v}{\Delta t}$$

$$\text{equations of motion} \quad v = u + at \quad s = \frac{(u+v)t}{2}$$

$$v^2 = u^2 + 2as \quad s = ut + \frac{1}{2}at^2$$

$$\text{force} \quad F = ma$$

$$\text{work, energy and power} \quad W = F s \cos \theta$$

$$E_K = \frac{1}{2} m v^2 \quad \Delta E_P = mg\Delta h$$

$$P = \frac{\Delta W}{\Delta t}, P = Fv$$

$$\text{efficiency} = \frac{\text{useful output power}}{\text{input power}}$$

## MATERIALS

$$\text{density} \quad \rho = \frac{m}{V} \quad \text{Hooke's law} \quad F = k \Delta L$$

$$\text{Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} \quad \text{tensile stress} = \frac{F}{A}$$

$$\text{tensile strain} = \frac{\Delta L}{L}$$

$$\text{energy stored} \quad E = \frac{1}{2} F \Delta L$$

## WAVES

$$\text{wave speed} \quad c = f\lambda \quad \text{period} \quad T = \frac{1}{f}$$

$$\text{fringe spacing} \quad w = \frac{\lambda D}{s} \quad \text{diffraction grating} \quad d \sin \theta = n\lambda$$

$$\text{refractive index of a substance } s, \quad n = \frac{c}{c_s}$$

for two different substances of refractive indices  $n_1$  and  $n_2$ ,

$$\text{law of refraction} \quad n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$\text{critical angle} \quad \sin \theta_c = \frac{n_2}{n_1} \text{ for } n_1 > n_2$$